

Nematodes in *Sceloporus variabilis* (Squamata: Phrynosomatidae) in localities in the Altas Montañas region of Veracruz, Mexico

Mora-Collado, N.¹, Serna-Lagunes, R.¹, Tinoco-Camarillo, G.¹, Núñez-Pastrana, R.¹, Nogueta-Torres, B.², Salazar-Ortiz, J.^{3*}

¹Facultad de Ciencias Biológicas y Agropecuarias, Universidad Veracruzana, Veracruz México. nmora@uv.mx, gabytinoco1@hotmail.com, rserna@uv.x, rnunez@uv.mx. ²Laboratorio de Helmin-tología, Instituto Politécnico Nacional, bnogueta@ipn.mx. ³Colegio de Postgraduados, Campus Córdoba. Km. 348 Carretera Federal Córdoba-Veracruz, Congregación Manuel León, C. P. 94946, Amatlán de Los Reyes, Veracruz. México.

*Author for correspondence: salazar@colpos.mx

ABSTRACT

Objective: Identify nematodes of the digestive tract of the lizard *Sceloporus variabilis*.

Methodology: In different localities in the Altas Montañas region, 11 individuals of *S. variabilis* were collected and a range of morphological characteristics measured. Their intestines were dissected to identify the nematodes present.

Results: Were identified 17 nematodes belonging to the Oxyuridae family, present in the lizard *S. variabilis*. Not found differences in the frequency of nematodes between sexes and localities.


Implications: Identification of nematodes in *S. variabilis* is basic information for the knowledge of the parasite-host relationship.

Conclusions: The nematodes of the Oxyuridae family may cause mechanical symptoms in *S. variabilis*.

Key words: Reptilia, Squamata, Sauria, Phrynosomatidae, parasitism, host, Oxyuridae.

Diseases and their agents have been little studied in reptiles but some groups, such as the lacertilia, act as reservoirs, mainly of protozoa (Mata, 2018). Studying the influence of parasitic helminths on the life cycle and ecological processes of their hosts, is important in the context of describing parasite-host relationships (Aho, 1990). In addition, identifying endoparasites in reptiles is important to relating aspects of the biology of the species to their environment (Wilson and Carpenter, 1996).

In particular, the helminths in the pink-bellied spiny lizard (*Sceloporus variabilis* Wiegmann, 1834; Sauria: Phrynosomatidae) have not been identified. For this reason, the present study contributes to the knowledge of the diversity of nematodes present in the digestive tract of *Sceloporus variabilis*, a species of wide distribution on the American continent that is tolerant to disturbance of the ecosystems and has adapted to fragmented areas, presenting abundant populations (Chaves *et al.*, 2013). The genus *Sceloporus* is considered a biological model with which to study the parasite-host interaction, for which reason it is important to evaluate its role as a transmitter of nematodes towards



other groups of reptiles (Garduño-Montes de Oca *et al.*, 2017).

This study was conducted in three localities in the year 2019 (Figure 1): 1) Peñuela (18° 46' and 18° 58' N; 96° 49' and 96° 58' W); 2) Huiloapan de Cuauhtémoc (18° 48' and 18° 50' N; 97° 06' and 97° 10' W); and Palma Sola (18° 45' N and 96° 47' W). Eleven specimens of *S. variabilis* were collected and transported to the Laboratorio de Microscopía Óptica of the Facultad de Ciencias Biológicas y Agropecuarias, Orizaba-Córdoba region, of the Universidad Veracruzana and sacrificed through hypothermia induced by freezing (García, 2013). The sex of the lizards was determined by sexual dimorphism and the age category estimated from the measurement snout-vent length (SVL) < 42.4 mm for juvenile and > 42.4 mm for adults. A range of morphological characteristics was also recorded in the lizards (Table 1; Serna-Lagunes, 2005). The contents of the digestive tracts of the lizards were prepared in an aqueous solution in Petri dishes and observed under a microscope in order to quantify the frequency of nematodes per age category and sex of the lizard. An ocular micrometer was used to obtain the anatomical characteristics of the nematodes and these

were compared to examples in the scientific literature, books, laboratory manuals and specialized keys for the taxonomic identification of the nematodes (Anderson *et al.*, 1989).

The *S. variabilis* lizards evaluated in this study included three females (27% of the total) and eight males (73%). Regarding age category, 10 were adults (90% of the total) and one was juvenile (10%). The lizards presented a higher quantity of nematodes in Palma Sola (n=10) and Peñuela (n=7). In this sense, 17 morphotypes of nematodes were identified as belonging to the family Oxyuridae (Figure 2); four specimens [three males (75% of the total) and a female (25%)] of *S. variabilis* presented the highest prevalence of helminths (Table 1), while 36% of the lizards presented no nematodes, for which reason it is considered that *S. variabilis* does not act as a reservoir of nematodes. No association was presented between the nematodes and the sex, age category and localities of the lizards (X^2 ; $P > 0.05$).

On comparison of the results with those of Breves *et al.* (2011), studying Iguana iguana in Brazil, three males and three females presented more than 100 endoparasites

belonging to the family Oxyuridae. This family coincides with that reported in this study for *S. variabilis*. Salizar (2008) studied parasitic helminths of the Peru desert tegu (*Dicrodon guttulatum* Dumeril and Bibron, 1893) in Peru and, of 38 hosts sampled, 34 (89.47%) were parasitized with 7929 (99.85%) nematodes, with the males presenting greater parasitism (75.76%) than the females (15%). Comparing this with the case of *S. variabilis*, there was no significant frequency between sexes, which could be due to the difference in the number of hosts collected in each study.

On the other hand, Stephen *et al.* (2003) found 404 endoparasites (seven nematodes) in seven lizard species of the genus *Sceloporus* (including *S. variabilis*) of Mexico, a frequency




Figure 1. Sites at which the prevalence of nematodes was determined in *S. variabilis*. The map shows the sample size (n) at each sampling location.

Table 1. Morphometric variables of the pink-bellied spiny lizard *S. variabilis* and prevalence of nematodes (counts of nematodes present per individual).

No. Ind.	Male of <i>S. variabilis</i>	Total length (TL) cm	Snout-vent length (SVL) cm	Head width (HW) cm	Head length (HL) cm	Left leg length (LL)	Prevalence /absence of nematodes
1	Adult	13.5	6	1	2.1	1.7	0
4	Adult	9	4.5	1.5	2	0.5	5
5	Adult	13.2	5	1.2	2	2.7	0
6	Adult	13	4.5	1.5	3	2	0
7	Adult	9	4.5	0.5	1.5	3	3
8	Adult	15.5	7	1.7	1.5	3.5	0
9	Juvenile	4.5	2.5	0.5	0.8	1	0
10	Adult	12.5	5	1	1.5	1.5	0
	Mean	11.28	4.88	1.11	1.80	1.99	
	Standard deviation	3.30	1.22	0.42	0.60	0.96	
No. Ind.	Female of <i>S. variabilis</i>	Total length (TL) in cm	Snout-vent length (SVL) cm	Head width (HW) cm	Head length (HL) cm	Left leg length (LL)	
2	Adult	10	7	1.5	2	1.5	0
3	Adult	11	5.5	1	1.5	2	5
11	Adult	12	5.5	1	1.5	3	0
	Mean	11	6	1.16	1.66	2.16	
	Standard deviation	0.81	0.7	0.23	0.23	0.62	

of nematodes that was similar to that reported in this study. For this reason, the analysis of *S. variabilis* conducted in this study confirms the presence of nematodes of the family Oxysuridae, but this presence is not associated with the sex of the lizards.

ACKNOWLEDGEMENTS

Thanks go to the Laboratorio de Helminología, Instituto Politécnico Nacional, for their help with helminth identification, and to the project "Caracterización de recursos zoogenéticos de las Altas Montañas, Veracruz: aplicación de la filogeografía y modelación ecológica" (PRODEP: 511-6/18-9245/PTC-896) for funding and technical support of the study. Thanks also go to the Secretaria de Medio Ambiente y Recursos Naturales (SEMARNAT) for granting a scientific collection license for the purposes of teaching: SGPA/DGVS/001894/18 allowing collection of the studied specimens.

LITERATURE CITED

- Aho, J. M. (1990). Helminth communities of amphibians and reptiles: comparative approaches to understanding patterns and processes. In G. W. Esch, A. O. Bush y J. M. Aho. Parasite communities: Patterns and processes. (pp. 157-195). New York: Chapman y Hall.
- Anderson, R., A. Chabaud, and S. Willmott. (1989). Keys to the nematode parasites of vertebrates. Wallingford: CAB International Institute of Parasitology.
- Breves, P., M., Porto, A. Pissinatti, D. Luz & R.C. Menezes. 2011. Helminths oxiuridae parasitos de *Iguana iguana* (Squamata, Lacertilia, Iguanidae) procedentes do Brasil. Arquivo Brasileiro de Medicina Veterinária e Zootecnia, 63(6): 1574-1578.
- Chaves, G., Lamar, W., Porras, L.W. & Sunyer, J. (2013). *Sceloporus variabilis*. The IUCN Red List of Threatened Species 2013: e.T198414A2525951. <http://dx.doi.org/10.2305/IUCN.UK.2013-2.RLTS.T198414A2525951.en>. Downloaded on 19 August 2019.
- Garduño-Montes de Oca, E. U., J. D. López-Caballero & R. Mata-López. 2017. New records of helminths of *Sceloporus pyrocephalus* Cope (Squamata, Phrynosomatidae) from Guerrero and Michoacán, Mexico, with the description of a new species of Thubunaea Seurat, 1914 (Nematoda, Physalopteridae). ZooKeys 716: 43–62.
- García, M. R. A. 2013. Aspectos ecológicos y dimorfismo sexual de la lagartija vivípara *Sceloporus formosus* (Squamata: Phrynosomatidae) en la parte centro-este del estado de Puebla. Tesis de Licenciatura en Biología, Facultad de Estudios Superiores Zaragoza, Universidad Nacional Autónoma de México.
- Mata, R. 2018. Anfibios y Reptiles: enfermedades, hospederos y vectores. Departamento de biología evolutiva, Facultad de Ciencias, UNAM. México. Revista Latinoamericana de Herpetología, Vol.01 No.01.
- Salizar, V. P. 2008. Helminths parasites of *Dicrodon guttulatum*, Dumerily Bibron, 1893 (SAURIA: TEIIDAE) de la costa del Perú. Tesis de Biólogo con mención en zoología. Facultad de Ciencias Biológicas, Universidad Nacional Mayor de San Marcos. Lima Perú.













No.	Nematodes observed under a phase contrast microscope at magnification 400 X	
1	 <p data-bbox="552 442 738 472">Head of nematode</p>	 <p data-bbox="998 442 1169 472">Tail of nematode</p>
Nematodes observed under a compound microscope		
2	 <p data-bbox="519 804 771 834">Head of nematode (100X)</p>	 <p data-bbox="909 804 1258 834">Complete body of nematode (100X)</p>
3	 <p data-bbox="527 1123 763 1153">View of the head (100X)</p>	 <p data-bbox="982 1123 1193 1153">View of the tail (100X)</p>
4	 <p data-bbox="495 1442 795 1472">Observation of the head (400X)</p>	 <p data-bbox="941 1442 1226 1472">Observation of the tail (400X)</p>
5	 <p data-bbox="495 1740 795 1770">Observation of the head (400X)</p>	 <p data-bbox="941 1740 1226 1770">Observation of the tail (400X)</p>
6	 <p data-bbox="495 1942 795 1972">Observation of the head (100X)</p>	 <p data-bbox="941 1942 1226 1972">Observation of the tail (100X)</p>

Figure 2. Specimens of nematodes of the family Oxyuridae in the intestine of *S. variabilis*. A): Head. B): Tail. Observation under a microscope at 100X and 400X.

- Serna-Lagunes, R. 2005. Demografía de una población de *Sceloporus variabilis* en la localidad de Omiquila, municipio de Rafael delgado (Parque Nacional Cañón de Río Blanco), Veracruz. Tesis de Licenciado en Biología. Facultad de Ciencias Biológicas y Agropecuarias, Universidad Veracruzana. Córdoba.
- Stephen R., R. Charles, & R. Camarillo 2003. Gastrointestinal Helminths of seven species of Sceloporine lizards from Mexico. *The Southwestern Naturalist*, 48(2): 208–217.
- Wilson, S. C., & Carpenter, J. W. (1996). Endoparasitic diseases of reptiles. In *Seminars in Avian and Exotic Pet Medicine*. 2(5): 64-74.

